

NET WT:

18 Ibs (8.2 kg)



ORATORIES, INC.

KILLS RATS, MICE & MEADOW VOLES*

Kills Warfarin Resistant Norway Rats

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panels for First Aid and additional precautionary statements.

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-Ihroated woodrat, Southern plains woodrat, and Mexican woodrat

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

IF SWALLOWED:

Call a poison control center, doctor, or 1-877-854-2494, or 1-800-858-7378** immediately for treatment advice.
 Have person sip a glass of water if able to swallow.

FIRST AID

- · Do not induce vomiting unless told to do so by the poison control center or doctor.
- IF ON SKIN OR CLOTHING:
- Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- · Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
- ** Also call this number for information on health concerns and pesticide incidents.

NOTE TO PHYSICIAN

If swallowed or absorbed through the skin, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K₁ intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING If animal eats bait, call veterinarian at once. NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K₁. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transtusions may be needed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for sites, pests and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure

1. Store unused product out of reach of children and pets.

- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If ban can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, reaccours, bears, or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

Bait stations are mandatory for outdoor, above-ground user Tamper-resistant bait stations must be used wherever children, pets non-target mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This boddet may only be used to control the following rodent pests in and around man-made structures. House mouse (Mus musculus), Norway rat (Rattus naruegrous), Roof rat (Rattus raitus), Cotton mouse (Reformscus gossyninus), Cotton rat "Gigmodon hispidus), Deer mouse (Reromyscus maniculatus), Eastern harvest mouse" (Retirtodentemys humuii), Cotte mouse (Ochrotomys nutralii), Polynesian rat" (Rattus exulans), Meadow vole (Microtus pearsylvanius), Uniterfooded mouse (Peromyscus leucopus), White-Ihroated woodrat" (Neotoma albigula), Southern plains yoontat" (Neotoma micropus), and Mexican woodrat" (Neotoma mexicana). This product must be used in and within 100 feel of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to participant and estructures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to participant and structures to constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to participant provident interstations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied diredly to food or feed crops.



KILLS RATS, MICE, AND MEADOW VOLES*

Kills Warfarin Resistant Norway Rats

Norway rats, roof rats, and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

ACTIVE INGREDIENT:

 Bromadiolone (CAS #28772-56-7):
 0.005%

 OTHER INGREDIENTS†:
 09.995%

 †Contains Denatonium Benzoata
 TOTAL

 100.000%
 TOTAL

KEEP OUT OF REACH OF CHILDREN

See side panels for First Aid and additional precautionary statements.

STORAGE AND DISPOSAL

net contaminate water, food or feed by storage or disposal.

Pesticide Storage Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Wastes resulting from the use of this product may be placed in trash or delivered to an

approved waste disposal facility. Container Handling: Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling

or reconditioning; or puncture and dispose of in a sanitary landfil; or by incineration. In most states, burning is not allowed.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET WEIGHT: 18 lbs (8.2 kg)

EPA REG. NO. 12455-79 EPA EST. NOs. 12455-WI-1k,12455-WI-2^p, 12455-WI-3^w Superscript is first letter of the lot number.



DIRECTIONS FOR USE (Continued from other panel)

Burrow baiting with CONTRAC® All-Weather Blox is prohibited.

Do not place near or inside ventilation duct openings. Do per contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces viait dome into direct pontact with look. When used ju dSDA inspected facilities, this product must be applied in tamper-resistant bait startions. Bo not broadcast bait. Don't use this product in severs.

Do not sell this product in individual containers holding less than 16 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice, permeadow voles' will most likely find and consume the topic. Senerally, these aheas here along walls, by gnawed openings, in corners and concealed places, between fights and walls or in locations where lats, mice, or meadow voles', or their signs have been seen. Protect bait from rain and show. Rencyce as much alternique locat as possible.

APPLICATION DIFFECTIONS: RATS: Place 3 to 16 bait blocks (at intervals of 15 to 30 feet) per placement in mested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease. MICE AND MEADOW VOLES': Place 1 block per placement. Space placements at 8- to 12-too tintervals in infested areas. Two blocks

placements at 8- to 12-foot intervals in infested areas. Two blocks may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole" activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin or on clothing.

All handlers (including applicators) must wear: shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds and other wildlife. Dogs and predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

Product Code: CB4051

587CB-10